

**HCN3 Antibody**  
**HCN3 Antibody, Clone S141-28**  
**Catalog # ASM10184****Specification**

---

**HCN3 Antibody - Product Information**

Application	IHC, WB
Primary Accession	<a href="#">O88705</a>
Other Accession	<a href="#">NP_032253.1</a>
Host	Mouse
Isotype	IgG1
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Format	FITC

**Description**

Mouse Anti-Mouse HCN3 Monoclonal IgG1

**Target/Specificity**

Detects ~90kDa. No cross-reactivity against other HCNs.

**Other Names**

KIAA1535 Antibody, potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 3 Antibody, hyperpolarization activated cyclic nucleotide-gated potassium channel 3 Antibody

**Immunogen**

Fusion protein amino acids 660-779 (C terminus) of mouse HCN3

**Purification**

Protein G Purified

Storage **-20°C****Storage Buffer**

PBS pH7.4, 50% glycerol, 0.09% sodium azide

Shipping Temperature

**Blue Ice or 4°C****Certificate of Analysis**

1 µg/ml of SMC-306 was sufficient for detection of HCN3 in 10 µg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

**Cellular Localization**

Membrane


**HCN3 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## HCN3 Antibody - Images



Immunohistochemistry analysis using Mouse Anti-HCN3 Monoclonal Antibody, Clone S141-28 (ASM10184). Tissue: brain tissue. Species: Mouse. Fixation: 10% Formalin Solution for 12-24 hours at RT. Primary Antibody: Mouse Anti-HCN3 Monoclonal Antibody (ASM10184) at 1:1000 for 1 hour at RT. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 250-500  $\mu$ l for 5 minutes at RT. Localization: Granular staining of some Purkinje cells. Staining appears to be focal.



Western Blot analysis of Rat brain membrane lysate showing detection of HCN3 protein using Mouse Anti-HCN3 Monoclonal Antibody, Clone S141-28 (ASM10184). Load: 15  $\mu$ g. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-HCN3 Monoclonal Antibody (ASM10184) at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



Immunohistochemistry analysis using Mouse Anti-HCN3 Monoclonal Antibody, Clone S141-28 (ASM10184). Tissue: hippocampus. Species: Human. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-HCN3 Monoclonal Antibody (ASM10184) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.

### **HCN3 Antibody - Background**

Hyperpolarization-activated cation channels of the HCN gene family contribute to spontaneous rhythmic activity in both the heart and brain (1).

### **HCN3 Antibody - References**

1. Zong X., et al. (2005) J Biol Chem. 280(40): 34224-34233.